

BAVAFIT

early variety for crisps

NEW
RELEASES



BAVAFIT offers a very good adaptability to various environmental conditions. **BAVAFIT** is suitable for early deliveries to crisp factories.



QUALITY

dry matter content 23 – 24%
reducing sugar content very low
low sensitivity to mechanical damage
and internal defects

YIELD

medium number of tubers with uniform
size, medium–high marketable yield

RESISTANCES

- › PCN Ro 1, 4
- › Y-Virus (PVY) very high
- › potato leaf roll virus (PLRV) high
- › late blight medium
- › early blight medium
- › common scab medium to high

TUBER

oval, yellow skin, light yellow flesh

PLANT

white flowers, rapid development and
foliage cover

PLANTING

BAVAVIT can be planted on all types of
soil. BAVAVIT should be planted with
30 – 32 cm distance (in 75 cm rows) in a
well-prepared seedbed.
Protective chemical control against Rhi-
zoctonia solani is recommended.

FERTILISATION

up to 180 kg N / ha
minimum phosphorus 100 kg
potassium 250 – 300 kg / ha (preferably
sulphate)
sufficient magnesium 80 kg / ha.

In dry years foliage application with
micronutrients.

PLANT PROTECTION

BAVAVIT is not sensitive to Metribuzin,
routine chemical control of late blight
together with active ingredients against
early blight is recommended.

HARVEST / STORAGE

2 – 3 weeks after haulm killing skin set is
finished and BAVAVIT should be harves-
ted in time. Dormancy and storability is
very well.