

SANDRA

medium early, red skin

NEW
RELEASES



SANDRA shows high marketable yield, regular graded tubers with a nice red skin color even after long term storage.



QUALITY

cooking type B, no discoloration after cooking, starch content 13 – 14 %, low sensitivity to mechanical damage, internal defects and black spots, deep red skin

YIELD

very high marketable yield, uniform grading

RESISTANCES

- › PCN Ro 1
- › Wart disease K 1, 2
(6, 18 little susceptible)
- › Y-Virus (PVY) very high
- › potato leaf roll virus (PLRV) medium
- › late blight very high
- › black leg very high
- › common scab high

TUBER

oval to long oval tubers, shallow eyes, red skin, yellow flesh

PLANT

tall, very strong, flower red-violet
rapid development and foliage cover

PLANTING

SANDRA can be produced on all types of soil. Seed treatment against rhizoctonia is recommended. Normal planting depth and distance in the row, relatively late tuberisation

FERTILISATION

N-supply 100 – 120 kg N is recommended. Phosphorus and potassium according to local standard application

PLANT PROTECTION

no metribuzin sensitivity, due to a good tolerance to late blight spraying intervals can be less than average, haulm killing by splitting technique

HARVEST / STORAGE

very good storability, lower sensitive to *Helminthosporium solani* than comparable varieties